

Light Spectrum and Decay Constants in Full QCD with Wilson Fermions*

SESAM-Collaboration: U. Glässner^a, S. Güsken^a, H. Hoerber^b, Th. Lippert^b, G. Ritzenhöfer^b, K. Schilling^{a,b}, G. Siegert^b and A. Spitz^a.

^aPhysics Department, University of Wuppertal, D-42097 Wuppertal, Germany,

^bHLRZ c/o Forschungszentrum Jülich, D-52425 Jülich, and DESY, D-22603 Hamburg, Germany.

We present results from an analysis of the light spectrum and the decay constants f_π and f_V^{-1} in Full QCD with $n_f = 2$ Wilson fermions at a coupling of $\beta = 5.6$ on a $16^3 \times 32$ lattice.

1. INTRODUCTION

This analysis was performed as part of the SESAM project [1] to investigate **Sea** quark **E**ffects on **S**pectrum **A**nd **M**atrix-elements. Configurations were generated using the Hybrid-Monte-Carlo algorithm with the standard Wilson action at a coupling of $\beta = 5.6$ on a $16^3 \times 32$ lattice. We work at three different values of the dynamical quark mass: $\kappa_{\text{sea}} = 0.1560, 0.1570$ and 0.1575 corresponding to m_π/m_ρ -ratios of $0.83(1)$, $0.76(1)$ and $0.71(2)$ [2]. Our lightest quark-mass is approximately equal to the strange quark-mass. Up to now we generated about 8000 trajectories of unit-length with a time-step of $\Delta t = 0.01$. After measuring the integrated autocorrelation time for the plaquette and the pion-correlator at fixed timeslice [2,3] we decided to use configurations separated by 25 trajectories for the present spectrum analysis. Our sample consists of 100, 160 and 100 configurations for our 3 sea-quark values. Since we aim at a final sample of 3×200 configurations this talk should be considered as a half-time status report.

Quark propagators for the set of valence κ -values $\{0.1555, 0.1560, 0.1565, 0.1570, 0.1575\}$ were computed using the standard overrelaxed Minimal Residual Algorithm with smeared sources and local as well as smeared sinks. We used the Wuppertal gauge-invariant gaussian smearing method with $N = 50$ iterations and $\alpha = 4$, fitting the hadron-correlators to single-exponential functions. Throughout this analysis

we neglect correlations in time and in κ . We hope to present a stable correlated analysis on our final sample.

2. CHIRAL EXTRAPOLATIONS WITH FIXED SEA-QUARK MASS

To determine the critical hopping-parameter κ_c at a given value of κ_{sea} we fit the pseudoscalar masses according to

$$M_{PS}^2 = bm_q + cm_q^2 \quad (1)$$

with $m_q = 1/2(1/\kappa - 1/\kappa_c)$ being proportional to the quark mass. We find c to be significantly different from zero: a linear fit leads to a decrease of κ_c by four standard deviations with respect to the quadratic fit. The resulting values of κ_c are shown in table 1 together with their statistical errors. To estimate systematic errors on κ_c , we exclude the largest quark-mass from our data and find a change in κ_c of the order of one standard deviation.

For the chiral extrapolation of vector masses we use two different parametrizations

$$M_V = m_\rho + bm_q + cm_q^2 \quad (2)$$

$$M_V = m_\rho + bm_q + cm_q^{3/2}, \quad (3)$$

which are motivated by chiral perturbation theory. In contrast to the Π extrapolation we find that the values obtained from 2- and 3-parameter fits agree within errors; moreover eqs. 2 and 3 yield identical results. In table 1 we quote the values for the ρ -masses and the corresponding lattice-spacings as obtained from the

*talk presented by U. Glässner

κ_{sea}	κ_C	aM_ρ	a_ρ^{-1} [GeV]	M_N [GeV]	M_Δ [GeV]	f_π [MeV]	f_V^{-1}
0.1560	0.16065(8)	0.359(8)	2.14(5)	1.09(7)	1.30(9)	125(9)	0.33(3)
0.1570	0.15987(6)	0.341(8)	2.23(7)	1.14(6)	1.40(10)	101(8)	0.35(2)
0.1575	0.15963(11)	0.316(10)	2.44(8)	1.09(10)	1.22(8)	118(15)	0.30(3)

Table 1

Extrapolation results with fixed sea-quark mass. Errors quoted are purely statistical




Figure 1. *Meson extrapolations with fixed sea-quark mass.*

2-parameter-fits. The masses are calculated at κ_c because we find the mass-shifts induced by $\Delta\kappa = \kappa_c - \kappa_{\text{light}}$ to be negligible compared to the statistical errors². We will discuss the light quark masses estimated from $\Delta\kappa$ in detail in a forthcoming publication [2]. Figure 1 illustrates the variation of m_ρ with m_q and κ_{sea} .

The nucleon data (see figure 2) cannot be fitted with $c = 0$, but ansätze eqs. 2 and 3 do equally well. The quadratic extrapolation to κ_c leads to the values M_N quoted in table 1 together with the values for M_Δ . With the present statistics we do not see any sea-quark dependence of the nucleon mass in physical units.




Figure 2. *Nucleon extrapolations with fixed sea-quark mass.*

²In fact κ_c and κ_{light} still agree within errors




Figure 3. Pseudoscalar and Vector decay constants with fixed sea-quark mass.

We now consider the decay-constants, f_π and f_V^{-1} , defined as

$$\langle 0|A|\pi\rangle Z_k Z_A = f_\pi m_\pi \quad (4)$$

$$\langle 0|V_i|\rho\rangle Z_k Z_V = \epsilon_i f_V^{-1} m_\rho^2. \quad (5)$$

A and V_i are the local axial and vector currents on the lattice and Z_k , Z_A and Z_V the renormalization constants which connect the expectation values of equation 4 and 5 to the continuum. We determine these renormalization constants using tadpole improved perturbation theory [2,4]. The m_q -dependence of the decay-constants is presented in figure 3. Obviously the statistical accuracy is not yet high enough to resolve a κ_{sea} -dependence of f_π and f_V^{-1} at the chiral point (see also table 1).

3. EXTRAPOLATIONS IN κ_{sea}

A consistent method to extrapolate to zero sea-quark mass is to use only the data-points with $\kappa_{\text{sea}} = \kappa_{\text{val}}$ for the extrapolation. In the case of dimensionful observables one might argue about the impact of the varying scale along the trajectory. For this reason we consider only mass-ratios as shown in figure 4. We use linear fits only.




Figure 4. Extrapolations of mass ratios with $\kappa_{\text{sea}} = \kappa_{\text{val}}$.

M_N/M_ρ	1.6(2)
M_Δ/M_ρ	1.8(2)
f_π/M_ρ	0.14(3)
f_V^{-1}	0.32(6)

Table 2

Results from extrapolations in κ_{sea} .

The extrapolated values, quoted in table 2, agree (within large errorbars) with those obtained from extrapolations at fixed κ_{sea} .

4. CONCLUSIONS AND OUTLOOK

On our present statistics, uncorrelated data analyses of t - and κ_{val} distributions allow for stable chiral extrapolations in valence quark mass, m_q . We find the ρ -meson mass to be consistent with a linear chiral ansatz, while the remaining quantities considered exhibit *nonlinear* behaviour in m_q . m_ρ varies on the level of 15 % across our range of sea quark masses. Baryon masses and decay constants so far indicate a trend but produce no compelling evidence for such variation.

Chiral extrapolations of our present data in κ_{sea} itself still carry too large uncertainties to be reliable. Better statistics and deeper penetration into the chiral regime are needed to improve on this situation.

REFERENCES

1. SESAM-Collaboration, Nucl. Phys. B (Proc. Suppl.) **47** (1996) 386.
2. SESAM-Collaboration, in preparation
3. SESAM-Collaboration, Preprint HLRZ 96-20 and WUB 96-17, to be published in Phys. Lett. B .
4. G.P. Lepage and P.B. Mackenzie, Phys. Rev. D **48** (1993) 2250.